

Notes on Hidden Mirror World

S. I. Blinnikov^{1,2,*}

¹*Institute for Theoretical and Experimental Physics (ITEP), 117218, Moscow, Russia*

²*IPMU, University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa, 277-8568, Japan*

A few remarks on Dark Matter (DM) models are presented. An example is Mirror Matter which is the oldest but still viable DM candidate, perhaps not in the purest form. It can serve as a test-bench for other analogous DM models, since the properties of macroscopic objects are quite firmly fixed for Mirror Matter. A pedagogical derivation of virial theorem is given and it is pointed out that concepts of virial velocity or virial temperature are misleading for some cases. It is shown that the limits on self-interaction cross-sections derived from observations of colliding clusters of galaxies are not real limits for individual particles if they form macroscopic bodies. The effect of the heating of interstellar medium by Mirror Matter compact stars is very weak but may be observable. The effect of neutron star heating by accretion of M-baryons may be negligible. Problems of MACHOs as Mirror Matter stars are touched upon.

1. INTRODUCTION

Dark Matter (DM) remains one of the main puzzles in modern physics and astrophysics. In this paper I have collected a few notes on one DM candidate, namely, Mirror Matter. I do not attempt to give a full review on the subject: only some comments on Mirror Matter properties from the point of view of astrophysics are put forward. The comments may be useful for particle physicists working on new models for DM, e.g. from hidden sector, because Mirror Matter serves as an example of the model, where not only particle properties, but also the properties of macroscopic objects are quite firmly fixed and well known.

* Electronic address: sergei.blinnikov@itep.ru

2. DARK MATTER MANIFESTATIONS

If Newton-Einstein’s law of gravity is not modified at large scales, then DM manifests itself in many circumstances.

First, in 1933, Fritz Zwicky has discovered virial paradox in Coma cluster of galaxies [1]: velocities of galaxies, u_{gal} , are on average much higher there than can be expected from an estimate of gravitational potential using only the mass of visible matter, M_{vis} :

$$u_{\text{gal}}^2 \gg u_{\text{vir}}^2 \sim \frac{G_N M_{\text{vis}}}{R}.$$

We will discuss below more accurate formulation of the virial theorem, and dangers of using notions like the “virial velocity” (u_{vir} in the above expression) in some situations. But here the discrepancy is real. Zwicky immediately suggested the presence of “dunkle Materie” [1], i.e. the Dark Matter to remove the discrepancy.

Very soon, just after the paper by Einstein [2] on gravitational lensing (already discussed by Chwolson [3] for ordinary stars), Zwicky has proposed that the presence of DM can be tested by the effect of gravitational lensing [4, 5]: this what we call now macrolensing. It is interesting to note that the idea was suggested to Zwicky by Vladimir Zworykin – the Father of television. (Zwicky writes in [4] that Zworykin has got the idea from R.W.Mandl, but I could not find a reference to any paper by R.W.Mandl on that.) Many wonderful examples of macrolensing are known now, like multiple images of distant quasars and galaxies, and they all show that the mass of the lens (an intervening cluster of galaxies) must be much larger than the mass of visible component in the lens (again, if general relativity gives the true description of gravity at relevant distances). The lensing on early stages of research was also discussed by Tikhov [6, 7]. Other tools, important especially for compact dark objects, like microlensing were developed later [8–14], see a review in [15].

Very important for DM studies is the dynamics (e.g. rotation curves) of galaxies, also first proposed by Zwicky in 1937 [16]. Real measurements of missing mass growing with radius in M31 (Andromeda) galaxy have been performed already at that time [17]. But Zwicky’s seminal ideas and papers were almost forgotten, and rediscovery of DM occurred only four decades later, in 1970s. Then it was much more popular in USSR, than in the West, where it was accepted only after strong resistance, see the review by Einasto [18].

New evidence in favor of DM came from discovery of huge amount of X-ray emitting gas in clusters of galaxies. Mass of this gas is order of magnitude larger than the total visible

mass of galaxies, but still not sufficient to resolve the virial paradox. The temperature of the gas, $T \sim$ a few keV, corresponds to the velocity of protons of the same order as relative velocities of galaxies u_{gal} in a cluster. The hot gas cannot be trapped by the gravity of visible baryons, and this also points to the need of DM.

More detailed history of DM discovery see in [18–21].

Probably, the most important manifestation of DM is its influence on structure formation. A homogeneous universe in Friedmann solutions contains no galaxies or clusters – no structures. Matter in Universe was initially almost uniform, with small perturbations left after inflation [22–29] (or another initial stage, e.g., in cyclic scenarios [30]). After the recombination gravity enhances the density contrast, leading to the formation of structures when the self-gravity of an over-dense region becomes large enough to decouple the blob of matter from the overall expansion. Cosmic Microwave Background (CMB) fluctuations tell us that perturbations in density of baryons at epoch of recombination at redshift $z = z_{\text{rec}} \sim 10^3$ were on the level of 10^{-5} . They can grow only as the scale-factor $a \propto (1 + z)$ according to the Lifshits theory of growth of small perturbations in an expanding universe. Thus, now, at $z = 0$, we would get not more than 1% fluctuations of density, contrary to the observed structure.

DM particles can have larger amplitude of density perturbations and they begin to form structures earlier, while the baryonic matter is overwhelmed by photon pressure until the time of recombination. If DM dominates in gravity, the “dark matter boost” is provided for structure formation in visible matter, see e.g. [31, 32].

3. DARK MATTER CANDIDATES AND MIRROR MATTER

Current estimates of the fraction of DM in the average density of Universe are around 25% (assuming standard gravity), while baryons cannot contribute more than a few percent as found from abundances of light elements in primordial nucleosynthesis [33] and analysis of CMB [34].

If Dark Matter is real and non-baryonic, the most acute question is: “What is DM made of”? A clear and consize introduction to possible DM candidates is foind in G.Raffelt’s reviews [35, 36]. From the last one, at “The Dark Universe” conference, Munich, May 2004, one can point out a few probable models for DM.

- Supersymmetric particles — neutralino for heavy WIMPs (weakly interacting massive particles), or gravitino for lighter ones and their clusters.
- Axion-like particles and objects made of those particles.
- Mirror Matter particles and objects [37]. This is not so popular now as a DM model, but actually it is the oldest proposed candidate.

In spite of years of intensive search for WIMPs, nothing yet is discovered in laboratory! That is why Mirror Matter $\equiv$ MM is still a viable Dark Matter candidate: MM particles in the simplest MM model interact with ordinary matter only gravitationally (if we do not involve weak electromagnetic coupling like in [38]), and they can not be detected in laboratory experiments.

Neutralino and axions have electromagnetic interactions (on weak scale). MM in pure form has no electromagnetic (or other standard model) interactions with Ordinary Matter $\equiv$ OM, only gravity. MM has analogs of all “our” interactions within Mirror sector. There are also models with slightly “broken” Mirror symmetry which have weak electromagnetic coupling to OM.

Hypothesis on existence of invisible (or Dark) Mirror Matter was put forward by Kobzarev, Okun and Pomeranchuk in 1966 in the paper “On the possibility of observing mirror particles” [37] after the discovery of P- (1956) and CP-parity non-conservation (1964), when the mirror asymmetry of visible matter had become obvious. They have considered possible consequences of existence of matter which is mirror relative to our matter.

The history is often distorted in modern reviews: Lee and Yang [39] introduced a concept of right-handed particles, but their “R-matter” was not hidden! – as some of recent papers claim, e.g. [40]. An exact citation from [39] reads:

“...the interaction between them is not necessarily weak. For example, p_R and p_L could interact with the same electromagnetic field and perhaps the same pion field. They could then be separately pair-produced, giving rise to interesting observational possibilities”.

Kobzarev, Okun and Pomeranchuk [37] have shown that common electromagnetic and strong interaction of O and M particles are not allowed, this would contradict experiments. They demonstrated that only gravity and very weak interaction were possible between the two kinds of particles. So, it was shown that Mirror Matter lies in the hidden sector for the first time in [37]. The hidden world of Mirror Matter has the same microphysics as the

“visible” one. The term “Mirror Matter” was introduced first time also in [37]: it was taken by the authors from the book by Lewis Carroll “Through the Looking-Glass”.

Another important paper by L.B. Okun, “On the search for new long-range forces”, [41, 42] discussed other DM forms. The idea of MM search by the effect of gravitational lensing was also put forward in [41, 42].

Our papers with M.Yu.Khlopov “On possible effects of ‘mirror’ particles” [43–47] began systematic studies of astrophysics of MM as a form of DM.

After [37] about 300 papers have been published (among them e.g. Z.Berezhiani, R.Foot were very active, e.g. [48–50]), an extensive review of the literature on the subject is written by Okun [51]. More recent (brief) review by Z.K. Silagadze has an intriguing title “Mirror dark matter discovered?” [52]. However, to remove the question mark in this statement would be premature: Mirror world is still hidden from us.

4. VIRIAL PARADOXES

The discovery of DM by Zwicky was based on virial theorem. Here I derive the virial theorem and show that for some DM candidates its application is not trivial. We have for energy E of a stationary state for a system with Hamiltonian H

$$\delta E_{\text{tot}} \equiv \delta \mathcal{E} = \delta \langle H \rangle = 0$$

in the first order of perturbation of its wave function $\delta\psi$, i.e. the variational principle in quantum case. This is a starting point for seeking for the estimates of the ground state energy of quantum systems using trial approximation functions for ψ in standard applications of the variational principle. More interesting for us is the virial theorem. Compare the text below with derivation of virial theorem by Fock [53] in a quantum system, here I present it for a particular N-body problem with Coulomb potential.

Let us take a perturbation of the form:

$$\psi + \delta\psi = \alpha^{3N/2} \psi(\{\alpha \vec{r}_i\}), \quad i = 1, \dots, N,$$

so the wave function is uniformly changed for all $3N$ space coordinates. The coefficient $\alpha^{3N/2}$ is from normalization to unity. Note, that $\alpha > 1$ here corresponds to the *compression* of the whole system.

Schrödinger equation:

$$i\hbar \frac{\partial \psi}{\partial t} = H\psi$$

with

$$H = - \sum_j \frac{\hbar^2}{2m_j} (\nabla^2 \psi) + U\psi$$

gives after averaging

$$\langle H \rangle = \langle E_{\text{kin}} \rangle + \langle U \rangle.$$

Now we have after the perturbation:

$$\langle E_{\text{kin}} \rangle \rightarrow \alpha^2 \langle E_{\text{kin}} \rangle$$

because for non-relativistic (NR) particles, $E_{\text{kin}} \propto p^2 \propto 1/\lambda^2$, i.e.

$$\frac{\partial^2 \psi(\dots \alpha x_i \dots)}{\partial x_i^2} = \alpha^2 \frac{\partial^2 \psi}{\partial x_i^2}.$$

The variation of U depends on the law of interaction. For Coulomb (and Newton!) interactions

$$\langle U \rangle \propto \int \sum_{i \neq k} \psi^* \frac{1}{r_{ik}} \psi d^N \vec{r} \rightarrow \alpha \langle U \rangle.$$

Thus

$$\langle H \rangle \rightarrow \alpha^2 \langle E_{\text{kin}} \rangle + \alpha \langle U \rangle,$$

and variation of this gives

$$\delta \langle H \rangle = 2\alpha \delta \alpha \langle E_{\text{kin}} \rangle + \delta \alpha \langle U \rangle = 0,$$

so with $\alpha = 1$ for unperturbed state we find

$$2\langle E_{\text{kin}} \rangle + \langle U \rangle = 0.$$

This is the virial theorem for atomic Coulomb potential and for globular stellar clusters, and for clusters of galaxies as well! See the use of Schrödinger equation for stellar dynamics in [54, 55].

For all those systems (NR atoms or plasma, NR stars and clusters)

$$\mathcal{E} = \langle E_{\text{kin}} \rangle + \langle U \rangle = -\langle E_{\text{kin}} \rangle,$$

so the loss of total energy $\mathcal{E}$ corresponds to the *growth* of the kinetic energy $\langle E_{\text{kin}} \rangle$. The same is true for the internal energy of matter if it is in the form of kinetic energy of particles.

We should remember that we derived this based on the stationary quantum state which corresponds to an orbit in a classical cluster. The cluster must be in a relaxed (so-called “virialized”) classical state, i.e. it should be considered on a time-scale much longer than the orbital period. Otherwise, e.g. on a stage of violent relaxation, the virial theorem cannot be applied.

Now let us consider *non-interacting* particles in a potential well. From Schrödinger equation for NR particles (and also for classical objects!), if $U \propto r^k$, we find in the same way as above:

$$2\langle E_{\text{kin}} \rangle - k\langle U \rangle = 0.$$

If $k = -1$ we have Coulomb and Newton laws, for $k = 2$ — a harmonic oscillator.

The case $k \rightarrow +\infty$ corresponds to a hard reflecting wall, then $\langle U \rangle$ tends to zero relative to $\langle E_{\text{kin}} \rangle$, but the variation of $\langle U \rangle$ is always of the same order as the variation of $\langle E_{\text{kin}} \rangle$. Since the ‘force’ on the particle is $-\nabla U$, we define pressure P (i.e. the ‘force’ on unit area, $A = 1$, of the wall) so as

$$\delta\langle U \rangle = PA\delta x = P\delta V.$$

For our variations $V \rightarrow V/\alpha^3$, and when we express variations of H again through α , we get

$$2\langle E_{\text{kin}} \rangle - 3PV = 0,$$

i.e. the same relation as in classical derivation when the moments of equation of motion are averaged in time (see e.g. Kubo’s textbook on statistical mechanics).

Thus, we have for pressure in NR case:

$$P = \frac{2}{3}\varepsilon$$

so

$$E_{\text{kin}} = E_{\text{thermal}} = \frac{3}{2} \int P dV .$$

Important for us is also an extremely relativistic (ER) case [53]: $E_{\text{kin}} \propto p \propto 1/\lambda$, then

$$\langle E_{\text{kin}} \rangle + \langle U \rangle = 0.$$

For ER case:

$$P = \frac{1}{3}\varepsilon$$

and

$$E_{\text{kin}} = E_{\text{thermal}} = 3 \int P dV .$$

and from virial theorem in both NR and ER cases we have got

$$3 \int P dV + \langle U \rangle = 0 .$$

Of course this relation is valid for arbitrary $P = P(\varepsilon)$ and its derivation can be found in standard textbooks.

Now let us discuss why it is dangerous sometimes to use the virial theorem and to rely on T_{vir} and v_{vir} for DM particles with long-range interactions.

For $U \sim -G_{\text{N}}M^2/R$, omitting all coefficients of order unity, we find that pressure and density in the center of a spherical cloud of mass M and radius R are:

$$P_c \simeq \frac{G_{\text{N}}M^2}{R^4}, \quad \rho_c \simeq \frac{M}{R^3}.$$

and we have

$$P_c \simeq G_{\text{N}}M^{\frac{2}{3}}\rho_c^{4/3}.$$

So, if we have a classical ideal plasma with $P = \mathcal{R}\rho T/\mu$, where $\mathcal{R}$ is the universal gas constant, and μ — mean molecular mass, we get

$$T_c \simeq \frac{G_{\text{N}}M^{2/3}\rho_c^{1/3}\mu}{\mathcal{R}}.$$

With $\mu \simeq 1$ for hydrogen-helium fully ionized plasma we get for the Sun $T_c \simeq 10^7 \text{ K} \simeq 1 \text{ keV}$. This is OK for “virial” velocity $v_{\text{vir}}^2 \sim \varphi \sim r_g/R_{\odot} \sim 10^{-6}$ (although $\sqrt{2}$ less than for neutrals!). The temperature T_c may be called a “virial” T_{vir} for ions, but not for electrons, the latter have much higher velocity than “virial”, since they have the same temperature but much smaller mass.

The discrepancy is much larger in degenerate stars. E.g. in Sirius B, with its central density higher than 10^7 g cm^{-3} , we have velocity of electrons $\sim c$ (due to high Fermi momenta in degenerate electron gas) and velocity of ions tends to zero. They are hot enough to form a classical ideal plasma, but T of course is quite different from the classical virial value.

This has direct relevance to DM studies. One can imagine a DM candidate like a neutrino of a few eV mass. It will determine the mass of a cluster or super-cluster of galaxies [56], but T_{vir} loses its sense when degeneracy becomes important. Situation with long-range

forces (like, e.g., in models for hidden charged Dark Matter [40]) is analogous with ordinary plasma, and neither virial velocity, nor virial temperature are appropriate concepts for DM candidates of this sort.

This paradoxical conclusion does not mean that the virial theorem is not valid. Simply its standard derivation, and the formula $3 \int P dV + \langle U_g \rangle = 0$, where U_g is the gravitational potential energy, mask the fact that all other types of potentials must be accounted for. Inside a star electrons always have velocity much higher than u_{vir} , either due to temperature which must be equal to that of ions, or due to high Fermi momentum in a cold star. So electrons tend to fly away from the gravitational potential well, and immediately a very small polarization of plasma occurs. Electrons feel electrostatic field which accelerates them much stronger than gravity because of their small mass. Ions feel electric force of the opposite sign which equilibrates the gravity. With account of the electrostatic potential, added to the gravitational potential in the virial theorem, the particles of different sign have velocities of the right order.

The concept of pressure P has to be used with care as well. In a degenerate dwarf the pressure of electrons is much higher than the pressure of ions, but it cannot be said in general that P describes the transfer of momentum in collisions of electrons and ions, like in classical fluids: there are no individual collisions for degenerate particles. Electrons cannot lose their momentum: the lower states are all occupied. They are in steady states (without collisions) in a potential well of electric field with a size of star. Ions may form an ideal gas, or liquid, or even a crystal and they are supported against the gravity by the *volume* force of the same electric field. So P enters here not like a *surface* force on a “brick wall”, but just as a flux of momentum per unit time mediated by electric field.

5. ACCRETION ONTO M-OBJECTS

Some popular DM candidates like neutralino are not quite “dark”. They may, e.g., annihilate and produce gamma-rays. Actually, this is one of the most promising way to identify them in space.

Pure MM manifests itself via gravity only. The only source of radiation of O-photons by an M-star is the accretion of the O-matter. The term “accretion” denotes here not only the capture of matter by a star, but the whole complex of the processes of gravitational




Figure 1. Possible regimes of accretion onto an M-object moving supersonically through O-matter from [43]:

- a) $R \ll R_A$; HS denotes a head (bow) shock, TS — a possible trailing shock; the matter is partially captured and cools down in the potential well of M-star.
- b) $R \approx R_A$; S is a shock front; the matter is virtually not captured.
- c) $R > (u/u_s)R_A$; WS denotes a weak shock (in general, presumably, may not form at all); the matter flow is only slightly perturbed, no capture.

interaction of the M-star with the surrounding O-medium. MM stars may accrete ordinary matter and one may hope to detect a signal of an unusual accreting object. Early proposals to search MM objects along these lines have been put forward in [43, 46, 47].

Standard accretion of gas on gravitating center is described by so called Bondi–Hoyle–Lyttleton (BHL) theory, see the review by R.G. Edgar [57] for references. Details of accretion of non-interacting particles on a gravitating center are considered in [58, 59].

According to the classical theory of accretion, the gravitating body of mass M effectively

captures the matter within the accretion radius

$$R_A = 2GM/u^2 ,$$

where u is the velocity of the body's motion through the medium in the supersonic case, or the sound speed in the medium, u_s , in the subsonic case. We may use very crude estimates here, so the sound speed is roughly related to the temperature by $kT = mu_s^2$. The square of R_A gives the cross-section $\propto \pi R_A^2$ and the result is essentially the same as for Rutherford scattering (replacing GM by Ze).

For $u > 10$ km/s the relative motion is supersonic, since $u_s \simeq 10$ km/s for the temperature of the medium $T \sim 10^4$ K. Supersonic accretion onto M-star was considered qualitatively by us [43, 46, 47]. If the radius of a star R is much smaller than R_A , then accretion must be similar to accretion onto a black hole or a neutron star but there is no trapping surface. For larger radii there are different regimes possible, see Figure 1, depending on ratio of the star radius R and the accretion radius R_A .

For a very large radius R there is hardly any hope to detect the emission of the perturbed gas. So the case $R \ll R_A$ which obtains easily for normal stars, and is especially good for white dwarfs and neutron stars, is worth to study. This case must be close to the BHL accretion.

For such a case G. S. Bisnovatyi-Kogan et al. [60] have found a self-similar solution for a gas with constant adiabatic exponent γ . This solution has some singularities and numerical results [57] show a different picture, more similar to our Figure 1a.

Using formulae for accretion rate one can estimate the amount of mixing of O- and M-matter. From

$$R_A = \frac{2GM}{u^2} \approx 3 \times 10^{14} \frac{M}{M_\odot} \frac{1}{u_6^2} \text{ cm},$$

we get R_A about tens a.u. for $M = M_\odot$. If $\rho \sim m_p n$ is the density of the surrounding medium then maximizing the classical expression for a rate of accretion we get

$$\dot{M} = \pi R_A^2 \rho u \sim 10^{12} \left(\frac{M}{M_\odot} \right)^2 \frac{n}{u_6^3} \text{ g/s}$$

and for $n \sim 1 \text{ cm}^{-3}$ a star accretes $3 \times 10^{28}/u_6^3$ g in 10^{10} yrs, a bit heavier than Earth mass.

Luminosity of infalling gas is approximately given by

$$L \sim \dot{M} \varphi_0 ,$$

where φ_0 is the characteristic value of the gravitational potential in the place of the stopping of the infalling matter. If the radius of an accreting M-star R is of order of the neutron star radius, then $R \ll R_A$, $\varphi_0 \sim 0.1c^2$ and L will approach a few percent of Solar luminosity for a single M-star. Luminosity may be much higher in a binary of O-M stars, however, the probability to find such a binary is very low [46, 47].

It is clear that to discover a MM star by its accretion of ordinary matter is extremely hard. It is hard even for a single old neutron star made of ordinary visible matter. Nevertheless, the chances are not absolute zero. Crude estimates of radiation fluxes produced by accretion onto a MM star (or any other hidden matter star) can be obtained as follows.

Particle number density can reach $n \sim 10^6 \text{ cm}^{-3}$ in molecular clouds. It can be higher in a binary in a young stellar cluster. Then for a MM neutron star one gets up to $L \sim 10^{38}$ ergs/s at $n \sim 10^6 \text{ cm}^{-3}$. For a MM white dwarf the luminosity $L \sim 10^{36}$ ergs/s, but the number of MM white dwarfs should be much larger.

It is hard to predict the spectral distribution without detailed computations. The hardest part of the spectrum should be determined by the stopping radius (if there is no strong non-thermal tail). For a neutron star it may go to X-rays. It might be softer for a white dwarf. But in both cases there is **no** solid surface for O-matter in the beginning of accretion. Later, after accumulation of some amount of ordinary matter, such a surface may build up, and spectrum will change.

The soft part of the spectrum may be determined by a total area of excited region and by coherent radio emission. For a black body and a surface of $4\pi R_A^2$ this implies $T_{\text{bb}} \sim$ a thousand K, and a near IR wave-range for the maximum for this high $L \sim 10^{38}$ ergs/s. But for $L \sim 4 \times 10^{31} \sim 0.1L_\odot$ the Wien peak is already in sub-mm range.

If we have a sensitivity of 3 mJy then to detect an object located at 100 pc we need it to have a luminosity not less than $L_\nu \sim 10^{16}$ ergs/(s · Hz). Thus, an instrument like RadioAstron may detect an accreting MM white dwarf if only $\sim 10^{-3}$ of its weak luminosity at $n \sim 1 \text{ cm}^{-3}$ ($\sim 0.001L_\odot$) goes into radio (for $\nu < 25 \text{ GHz}$). Or much smaller fraction: 10^{-10} — in case of $L \sim 10^{36}$ at $n \sim 10^6 \text{ cm}^{-3}$. This is quite plausible.

6. HEATING OF NEUTRON STARS BY MM

Now let us consider an opposite case: accretion of MM onto an O-star. It should have quite similar patterns of flow as in Figure 1 changing the flavours of particles accordingly. Effects of trapping MM by O-stars were discussed in our old papers [43, 46, 47].

An interesting effect is discussed in paper [61]: one can think that neutron stars capturing mirror matter could be heated at the order of 100 MeV per mirror baryon. The cause of the heating can be ascribed to the necessity to emit the excessive binding energy: after accretion the final state has more mirror baryons than the initial state, the number of ordinary baryons is constant, but the energy per ordinary baryon is lower. Effectively the O-baryons fall into a deeper potential well.

I show that the number of 100 MeV per mirror baryon in heating is a strong overestimate. During the process of accretion of M-baryons the matter in neutron star really goes down into a deeper potential well. But it is compressed due to slow accumulation of MM, and this process is purely adiabatic for O-matter and there is **no** heating in O-sector. All heating and emission and restoration of energy balances occur in mirror-sector. I propose several simple Gedankenexperiments illustrating that.

Let us have an O-neutron star and cover it with an ideally reflecting surface. Let this star be ice cold. Let us call the baryons in this star “blue” ones. Let “red” (ordinary matter) baryons fall slowly onto this star. Of course they will emit radiation, but let the accretion rate be slow enough for them to cool down to the same ice temperature. The reflecting surface between blue and red baryons does not allow the interiors to absorb any radiation. The total mass of “red matter” grows together with the gravitational binding energy, total energy of “blue matter” decreases, the blue interiors do compress indeed, but they remain ice cold!

Now replace the red baryons with mirror matter and we need no reflecting surface at all, because the radiation of mirror photons (which keeps all energy balances) is harmless for blue baryons. There is no effect of heating.

When one compresses gas it heats up - but only if it is already hot, $T > 0, S > 0$. If one compresses adiabatically then T grows, but $S = \text{const}$. If we lose heat, then T may still grow, but S goes down (like in classical stars). Here I always mean S per unit baryon.

But now we discuss the case of cold objects, $T = 0, S = 0$. Whichever strong compression,

T does not grow if $S = \text{const.}$ It remains zero. To increase T one must have a strong non-adiabatic process, because S must grow. Slow accretion of mirror baryons is an adiabatic process for O-neutron star, hence no growth of T whichever deep potential well develops. One must produce a shock wave, or something like this, to heat it up.

Let us consider a charged ball of mass 0.1 kg falling in terrestrial gravity and zero electric field $\mathbf{E} = 0$ into a potential well (a dielectric hole) from 1 m. It must emit around one J until it comes to rest and to a cold state at the bottom of the well. Now let us switch on $\mathbf{E} \neq 0$ of such a strength that the potential well is much deeper, say 100 J. Now, after the falling, the ball must emit 100 J. Finally, let us take the situation similar to an accreting neutron star: let us have the ball already at the bottom of the well in case of $\mathbf{E} = 0$ (this is our neutron star and zero admixture of MM) and then let us start increasing $\mathbf{E}$ slowly (adiabatically!) – this is the analog of slow accretion of MM particles and adiabatic change of gravity – the ball comes to a very deep potential well of 100 J. But it emits virtually nothing: it feels only adiabatic compression by electrostatic force, and if its temperature was $T = 0$ K initially, it will remain so forever! The same must be for the neutron star accreting MM.

If the reader does not like the electric field, one can do the same Gedankenexperiment in a potential well with artificial gravity (say in an accelerated laboratory, changing the force of engines adiabatically, i.e. slowly).

To make my counterexamples extreme let us take one O-atom, say of H and put it inside a cloud of rarefied M-matter. Let the M-cloud eventually to collapse to a star, and then to an M-neutron star with our atom in its center. It is still cold but it is in a deep potential well, like a 100 MeV/baryon solely due to the gravity of M-baryons. We need this order of energy to pull it out of the well. Do we need any process for our atom to emit 10 % of its mass? No, it could not do so. It was never excited. The energy was surely emitted by M-baryons in their M-photons, neutrinos etc. because they had to stop during accretion.

This is clear also in quantum language: any system in a ground state will stay in this state when the potential changes adiabatically. There are no excitations, no heating unless there are non-adiabatic perturbations.

That is why the Hawking radiation of black holes is so weak (if any): the adiabaticity of virtual modes can be violated only for wavelengths of order R_g , hence such a low T of a BH (due to Wien law for a wavelengths of a few km) for stellar masses.

The limits on neutron star heating by accretion of mirror baryons and other intersing

effects are discussed more quantitatively in the paper [61].

7. MICROLENSING

Invisible stars can be found by effect of microlensing. Those objects are called now MACHOs. Below I will write MACHO for “Massive Astrophysical Compact Halo Object” and MACHO for the collaboration [62] studying them. For mirror stars the effect of microlensing as a means to discover them was discussed in [63–65]. The difference with macrolensing is that the image is not resolved, the effect is in the enhancement of the detected flux of the lensed object due to the passage of the lens across the line of sight.

The MACHO results [62] of photometry on 11.9 million stars in the Large Magellanic Cloud (LMC) has revealed 13 - 17 microlensing events. This is significantly more than the ~ 2 to 4 events expected from lensing by known stellar populations. If true, this points to the existence of stellar-like invisible objects in halo of Milky Way, perhaps MM stars.

Not all DM in the halo may be in MACHO, only a fraction, usually denoted by f . MACHO group (LMC) has given for the halo fraction $0.08 < f < 0.50$ (95% CL) to MACHOs in the mass range $0.15M_{\odot} < M < 0.9M_{\odot}$. EROS collaboration has placed only an upper limit on the halo fraction, $f < 0.2$ (95% CL) for objects in this mass range. Later they have given $f < 0.1$ for $10^{-6}M_{\odot} < M < 1M_{\odot}$ (see refs. in [66]). AGAPE collaboration (working on microlensing of M31 - Andromeda galaxy) finds the halo fraction in the range $0.2 < f < 0.9$. (See [67] where arguments are given in favor of true halo MACHOs against self-lensing by ordinary stars in M31.) The MEGA collaboration marginally conflicts with them finding a halo fraction $f < 0.3$ [68].

Thus, the results of different groups are partly conflicting and quite confusing for a theorist. Some workers even declared the End of MACHO Era (1974-2004), e.g. [69].

Based on the results of MACHO group Bennett [70] has concluded that MACHOs are real and lensing optical depth is $\tau = (1.0 \pm 0.3) \times 10^{-7}$. This is not far away from the old MACHO value of $\tau = 1.2 \times 10^{-7}$, but the error is so large that it is within $\lesssim 2\sigma$ of the effect from ordinary stellar populations alone [71]. Evans and Belokurov [71] confirm lower number (optical depth τ) of MACHOs to the LMC from EROS [66] collaboration, who have reported $\tau < 0.36 \times 10^{-7}$. But there is a new paper of the same group – in favor of MACHOs from binary star studies [72] putting arguments against the pessimistic conclusions of [69].

Most recent study with HST (Hubble Space Telescope) follow-up shows that MACHO collaboration data are not contaminated by background events [73], and this adds some optimism to those who believe that invisible stars do exist.

Microlensing results do not allow us to have too many invisible stars. Let us take their local density like $8 \cdot 10^{-4} M_{\odot}/\text{pc}^3$ (solar masses per cubic parsec), i.e. *two orders of magnitude lower* than the number $0.076 \pm 0.015 M_{\odot}/\text{pc}^3$ given by Hipparchos satellite from accurate studies of dynamics of visible stars.

If about half of them are white dwarfs (this is quite likely in MM-sector) then we may have $\sim 2 \cdot 10^3$ mirror white dwarfs within 100 pc – just one order less than visible white dwarfs (Holberg et al. [74]) – and the chances to find a few of them very close to us appear nonzero.

8. LIMITS ON SELF-INTERACTING DM

Recently the cluster of galaxies 1E 0657-558 (“Bullet” cluster) has drawn general attention. It is an example of collision of two clusters, which gives new clues to the proof of reality of DM and pauses difficulties for alternatives to DM like modified gravity (MOND). Here it is clear that the DM distribution, which is traced by the effect of lensing, follows the distribution of stars and galaxies (which are effectively a collisionless gravitating gas) and is shifted from hot X-ray emitting gas which dominates the baryon mass in the cluster [75–77].

Many physicists tend to use this example as a direct proof of a small cross-section of self-interaction of DM particles. In reality it proves only what is observed and nothing more: DM behaves like ordinary stars which interact only by gravity and form a collisionless matter. But of course O-stars are made of strongly and electromagnetically interacting particles.

Let us take another example, the cluster MACS J0025.4-1222 which is similar to the Bullet cluster. It is found there that the DM cross-section obeys the limit $\sigma/m < 4 \text{ cm}^2 \text{g}^{-1}$ [78]. But this limit would be true for DM particles only if they do not form bound objects! If DM there is anything like our normal solid bodies, say ice with density like 1 g cm^{-3} with the size r larger than a few cm, then this limit is very well satisfied (σ is $\propto r^2$, while mass is $\propto r^3$).

So if the properties of DM particles are similar to our particles, and they are able to form stars, planets, asteroids, etc. (like MM), then all observations of the merging clusters

are reproduced. However, we have to squeeze a major fraction of MM particles into those compact objects, not leaving a large fraction in the form of gas like we have in OM in clusters of galaxies where the O-gas dominates in baryon component.

Dark matter in the Abell 520 cluster presents another case, difficult for CDM with WIMPs. In contrast to the Bullet cluster, the lensing signal and the X-ray emission coincide here, and are away from galaxies indicating that DM is collisional like in O-gas: "...a mass peak without galaxies cannot be easily explained within the current collisionless dark matter paradigm" [79]. Thus Abell 520 cluster indicates a significant self-interaction cross-section. In MM model we may get this leaving a major fraction of DM in form of gas. "It is hard for the WIMP based dark matter models to reconcile such a diverse behaviour. Mirror dark matter models, on the contrary, are more flexible and for them diverse behaviour of the dark matter is a natural expectation" [52].

L.B.Okun said: "MIRSY Dark Matter is richer than SUSY Dark Matter".

Other interesting examples, like Hoag's object, are also discussed by Silagadze [52].

9. CONCLUSIONS

I can draw a few conclusions from the above sketchy notes, anyway, some of them are important in general studies of Dark Matter candidates.

It is clear that the concepts of virial velocity or virial temperature are not always relevant.

One should not use the limits on self-interaction cross-sections derived from observations of colliding clusters as cross-sections for individual particles: they may form macroscopic bodies.

The effect of the heating of OM interstellar medium by MM neutron stars and white dwarfs is very weak but may be observable. One has to search for weak but high proper motion peculiar objects, especially in radio-range. Relation of these problems to RadioAstron mission and LOFAR [80] may be discussed and requires a detailed modeling. Peculiar binaries with low luminosity components, especially in star clusters, are also interesting.

The effect of the heating of OM neutron stars by accretion of M-baryons is negligible.

There are several arguments in favor for MM (or other hidden sector DM) search:

- Non-detection of WIMPs.

- Unknown properties of DM particles with respect to clustering on stellar-size and mass scale.
- Microlensing: lack of normal stars to explain MACHO events.
- Clusters of galaxies like Abell 520 which are hard to explain in pure CDM picture.
- New kinds of mysterious transients like reported by K. Barbary *et al.*, [81] may be explained by MM objects [61], anyway it must be a very unusual accretion event [82].
- One may speculate on possible MM life and intelligence (L.B.Okun, N.S.Kardashev).

MM-baryons may form stellar-mass/size compact objects, so they are interesting for microlensing, however, the current situation with observational data on MACHOs is not satisfactory: there are conflicting results of different groups. It seems that total DM halo mass in Galaxy cannot be explained by invisible stars, then more probable model for DM is a combination CDM+MM, or WDM+MM, where WDM is Warm Dark Matter. WDM may be, e.g., a sterile neutrino (with mass few keV) or gravitino [83–86].

Another window into the Mirror World may be photon - mirror photon mixing. There are strong limits on this mixing: orthopositronium experiments put an upper limit $\epsilon < 1.55 \cdot 10^{-7}$ (90% C.L.) [87]. Nevertheless, even smaller value of $\epsilon \sim 10^{-9}$ can explain DAMA/LIBRA experiments [38]. One should analyze experiments like DAMA, taking into account this option, and if $\epsilon \neq 0$ look for transformation of invisible mirror photons into visible ones. Other exotic effects are possible for $\epsilon \neq 0$, e.g. MM e^-e^+ transformations into OM e^-e^+ in M-pulsar, or a mirror magnetar, looking as a peculiar object with strong $B \sim 10^6$ G (D.Kompaneets).

I have not discussed cosmological evolution of MM. It is not clear a priori that pure MM can give sufficient amplitude of perturbations to provide the “DM boost” for visible matter structure formation: on the first glance, it seems that MM has a relatively short period of life in radiation-dominated era (contrary to WIMPs) when there is a logarithmic growth of perturbations. Nevertheless, MM is able to replace Cold Dark Matter (CDM) in evolution of perturbations [63, 88, 89]. The question was analyzed in detail in [88, 89] for various mixtures of MM and CDM. It is demonstrated [89] that the LSS (Large-Scale Structure) spectrum is sensitive to MM parameters, due to oscillations in mirror baryons and the collisional mirror Silk damping. If $T_{\text{MM}} < 0.3T_{\text{OM}}$ in Early Universe then M-baryons

will decouple from M-radiation before horizon crossing, and then as soon as they enter horizon, they do not oscillate, but grow exactly as CDM would do in radiation-dominated universe. Then CMB and LSS power spectra in linear regime are equivalent for mirror and CDM cases. For $T_{\text{MM}} > 0.3T_{\text{OM}}$ the LSS spectra strongly depend on the amount of mirror baryons, but for lower T_{MM} the entire dark matter could be made of mirror baryons given current observational limits on the CMB and LSS spectra.

Due to lower T_{MM} a larger fraction of mirror He is produced in cosmological nucleosynthesis, than in OM, hence faster stellar evolution follows [65]. It may result in lower fraction of MM intergalactic gas which is needed for explaining cases similar to Bullet cluster.

ACKNOWLEDGMENTS

This paper is written for a special issue of Yadernaya Fizika (Physics of Atomic Nuclei) dedicated to 80th birthday of L.B. Okun. I am very deeply grateful to Lev Borisovich for his lectures, books, joint work, and numerous discussions which always are a source of inspiration.

The paper is partly based on my talk at a conference “Radio Universe at Ultimate Angular Resolution” [90] and I am grateful to N.S.Kardashev for his invitation and encouragement. I thank M.I.Vysotsky, V.Lukash, D.Kompaneets, Z.Berezhiani, Z.Silagadze, A.Kusenko, P.Ciarcelluti, F.Sandin, for discussions and correspondence. My work is supported by IPMU, University of Tokyo, via World Premier International Research Center Initiative (WPI), MEXT, Japan. I am grateful to K.Nomoto and H.Murayama for their kind hospitality. The work in Russia is partly supported by the Russian Foundation for Basic Research grant RFBR 07-02-00830-a and by Russian Scientific School Foundation under grants NSh-2977.2008.2. NSh-3884.2008.2.

REFERENCES

-
1. F. Zwicky, *Helvetica Physica Acta* **6**, 110 (1933).
 2. A. Einstein, *Science* **84**, 506 (1936).
 3. O. Chwolson, *Astronomische Nachrichten* **221**, 329 (1924).
 4. F. Zwicky, *Physical Review* **51**, 290 (1937).
 5. F. Zwicky, *Physical Review* **51**, 679 (1937).
 6. G. A. Tikhov, *Dokl. Akad. Nauk. S.S.S.R.* **16**, 199 (1937).
 7. G. A. Tikhov, *Izvestiya Glavnoj Astronomicheskoy Observatorii v Pulkove* **16** (1938).
 8. S. Refsdal, *Mon.Not.Roy.Astron.Soc.* **128**, 295 (1964).
 9. S. Liebes, *Physical Review* **133**, 835 (1964).
 10. A. V. Byalko, *Astron.Zh.* **46**, 998 (1969).
 11. A. V. Byalko, *Soviet Astronomy* **13**, 784 (1970).
 12. A. V. Gurevich, K. P. Zybin, and V. A. Sirota, *Physics Letters A* **214**, 232 (1996).
 13. A. V. Gurevich, K. P. Zybin, and V. A. Sirota, *Uspekhi Fizicheskikh Nauk* **40**, 869 (1997).
 14. A. V. Gurevich, K. P. Zybin, and V. A. Sirota, *Physics Uspekhi* **40**, 869 (1997).
 15. A. F. Zakharov, *Physics of Particles and Nuclei* **39**, 1176 (2008).
 16. F. Zwicky, *Astrophys.J.* **86**, 217 (1937).
 17. H. W. Babcock, *Lick Observatory Bulletin* **19**, 41 (1939).
 18. J. Einasto, *ArXiv e-prints* (2009), 0901.0632.
 19. S. van den Bergh, *The Publications of the Astronomical Society of the Pacific* **111**, 657 (1999), [arXiv:astro-ph/9904251](#).
 20. A. Biviano, in *Constructing the Universe with Clusters of Galaxies* (2000), [arXiv:astro-ph/0010409](#).
 21. M. S. Roberts, in *Frontiers of Astrophysics: A Celebration of NRAO's 50th Anniversary*, edited by A. H. Bridle, J. J. Condon, and G. C. Hunt (2008), vol. 395 of *Astronomical Society of the Pacific Conference Series*, pp. 283–+.
 22. A. A. Starobinskii, *Soviet Journal of Experimental and Theoretical Physics Letters* **30**, 682 (1979).
 23. A. A. Starobinskij, *Physics Letters B* **91**, 99 (1980).

24. V. F. Mukhanov and G. V. Chibisov, Soviet Journal of Experimental and Theoretical Physics Letters **33**, 532 (1981).
25. K. Sato, Mon.Not.Roy.Astron.Soc. **195**, 467 (1981).
26. A. A. Starobinskij, Physics Letters B **117**, 175 (1982).
27. A. H. Guth and S.-Y. Pi, Physical Review Letters **49**, 1110 (1982).
28. A. A. Starobinskii, Soviet Astronomy Letters **9**, 302 (1983).
29. A. A. Starobinskii, Soviet Journal of Experimental and Theoretical Physics Letters **42**, 152 (1985).
30. B. Craps, T. Hertog, and N. Turok, ArXiv e-prints (2007), 0712.4180.
31. A. V. Gurevich and K. P. Zybin, Physics Uspekhi **38**, 2 (1995).
32. A. V. Gurevich and K. P. Zybin, Soviet Physics Uspekhi **38**, 687 (1995).
33. G. Steigman, Annual Review of Nuclear and Particle Science **57**, 463 (2007), 0712.1100.
34. D. N. Spergel, R. Bean, O. Doré, M. R. Nolta, C. L. Bennett, J. Dunkley, G. Hinshaw, N. Jarosik, E. Komatsu, L. Page, *et al.*, Astrophys.J. Suppl. **170**, 377 (2007), arXiv:astro-ph/0603449.
35. G. G. Raffelt, in *1997 European School of High-Energy Physics*, p. 235 - 278 (1998), pp. 235–278.
36. G. Raffelt, in *The Dark Universe in Munich* (2004), URL http://library.mppmu.mpg.de/webdocs/conf/confdoc/DUM04/Raffelt_DUM.pdf.gz.
37. I. Y. Kobsarev, L. B. Okun, and I. Y. Pomeranchuk, Sov. J. Nucl. Phys. **3**, 837 (1966), Yad. Fiz. **3**, 1154 (1966), in Russian.
38. R. Foot, Phys. Rev. D **78**, 043529 (2008), 0804.4518.
39. T. D. Lee and C. N. Yang, Physical Review **104**, 254 (1956).
40. J. L. Feng, H. Tu, and H.-B. Yu, Journal of Cosmology and Astro-Particle Physics **10**, 43 (2008), 0808.2318.
41. L. B. Okun, Zhurnal Eksperimentalnoi i Teoreticheskoi Fiziki **79**, 694 (1980).
42. L. B. Okun', Soviet Journal of Experimental and Theoretical Physics **52**, 351 (1980).
43. S. I. Blinnikov and M. Y. Khlopov, *Invisible objects inside the Sun and near the Solar system?*, Preprint ITEP-126 (1980).
44. S. I. Blinnikov and M. Y. Khlopov, Yad.Fiz. **36**, 809 (1982).
45. S. I. Blinnikov and M. Y. Khlopov, Sov. J. Nucl. Phys. **36**, 472 (1982).

46. S. I. Blinnikov and M. Y. Khlopov, *Astron.Zh.* **60**, 632 (1983).
47. S. I. Blinnikov and M. Y. Khlopov, *Soviet Astronomy* **27**, 371 (1983), *Astron.Zh.*, 60, 632-639.
48. Z. Berezhiani, D. Comelli, and F. L. Villante, *Physics Letters B* **503**, 362 (2001), arXiv:hep-ph/0008105.
49. Z. Berezhiani, P. Ciarcelluti, D. Comelli, and F. L. Villante, *International Journal of Modern Physics D* **14**, 107 (2005), arXiv:astro-ph/0312605.
50. R. Foot, *International Journal of Modern Physics D* **13**, 2161 (2004), arXiv:astro-ph/0407623.
51. L. B. Okun', *Physics Uspekhi* **50**, 380 (2007), arXiv:hep-ph/0606202.
52. Z. K. Silagadze, *ArXiv e-prints* (2008), 0808.2595.
53. V. Fock, *Zeitschrift fur Physik* **63**, 855 (1930).
54. L. M. Widrow and N. Kaiser, *Astrophys.J. Lett.* **416**, L71+ (1993).
55. R. Johnston, A. N. Lasenby, and M. P. Hobson, *ArXiv e-prints* (2009), 0904.0611.
56. G. S. Bisnovatyi-Kogan and I. D. Novikov, *Soviet Astronomy* **24**, 516 (1980).
57. R. Edgar, *New Astronomy Review* **48**, 843 (2004), arXiv:astro-ph/0406166.
58. A. F. Illarionov, *Astron.Zh.* **62**, 1080 (1985).
59. A. F. Illarionov, *Soviet Astronomy* **29**, 630 (1985).
60. G. S. Bisnovatyi-Kogan, Y. M. Kazhdan, A. A. Klypin, A. E. Lutskii, and N. I. Shakura, *Soviet Astronomy* **23**, 201 (1979).
61. F. Sandin and P. Ciarcelluti, *ArXiv e-prints* (2008), 0809.2942.
62. C. Alcock, R. A. Allsman, D. R. Alves, T. S. Axelrod, A. C. Becker, D. P. Bennett, K. H. Cook, N. Dalal, A. J. Drake, K. C. Freeman, *et al.*, *Astrophys.J.* **542**, 281 (2000), arXiv:astro-ph/0001272.
63. Z. G. Berezhiani, A. D. Dolgov, and R. N. Mohapatra, *Physics Letters B* **375**, 26 (1996), arXiv:hep-ph/9511221.
64. S. I. Blinnikov, *ArXiv Astrophysics e-prints* (1998), arXiv:astro-ph/9801015.
65. Z. Berezhiani, P. Ciarcelluti, S. Cassisi, and A. Pietrinferni, *Astroparticle Physics* **24**, 495 (2006), arXiv:astro-ph/0507153.
66. P. Tisserand, L. Le Guillou, C. Afonso, J. N. Albert, J. Andersen, R. Ansari, É. Aubourg, P. Bareyre, J. P. Beaulieu, X. Charlot, *et al.*, *Astron.Astrophys.* **469**, 387 (2007), arXiv:astro-ph/0607207.
67. A. Riffeser, S. Seitz, and R. Bender, *Astrophys.J.* **684**, 1093 (2008), 0805.0137.

68. G. Ingrosso, S. Calchi Novati, F. de Paolis, P. Jetzer, A. A. Nucita, G. Scarpetta, and F. Strafella, *Astron.Astrophys.* **462**, 895 (2007), arXiv:astro-ph/0610239.
69. J. Yoo, J. Chanamé, and A. Gould, *Astrophys.J.* **601**, 311 (2004), arXiv:astro-ph/0307437.
70. D. P. Bennett, *Astrophys.J.* **633**, 906 (2005), arXiv:astro-ph/0502354.
71. N. W. Evans and V. Belokurov, *Mon.Not.Roy.Astron.Soc.* **374**, 365 (2007).
72. D. P. Quinn, M. I. Wilkinson, M. J. Irwin, J. Marshall, A. Koch, and V. Belokurov, *ArXiv e-prints* (2009), 0903.1644.
73. C. A. Nelson, A. J. Drake, K. H. Cook, D. P. Bennett, P. Popowski, N. Dalal, S. Nikolaev, C. Alcock, T. S. Axelrod, A. C. Becker, *et al.*, *ArXiv e-prints* (2009), 0902.2213.
74. J. B. Holberg, E. M. Sion, T. Oswalt, G. P. McCook, S. Foran, and J. P. Subasavage, *Astron.J.* **135**, 1225 (2008).
75. D. Clowe, A. Gonzalez, and M. Markevitch, *Astrophys.J.* **604**, 596 (2004), arXiv:astro-ph/0312273.
76. D. Clowe, M. Bradač, A. H. Gonzalez, M. Markevitch, S. W. Randall, C. Jones, and D. Zaritsky, *Astrophys.J. Lett.* **648**, L109 (2006), arXiv:astro-ph/0608407.
77. M. Bradač, D. Clowe, A. H. Gonzalez, P. Marshall, W. Forman, C. Jones, M. Markevitch, S. Randall, T. Schrabback, and D. Zaritsky, *Astrophys.J.* **652**, 937 (2006), arXiv:astro-ph/0608408.
78. M. Bradač, S. W. Allen, T. Treu, H. Ebeling, R. Massey, R. G. Morris, A. von der Linden, and D. Applegate, *Astrophys.J.* **687**, 959 (2008), 0806.2320.
79. A. Mahdavi, H. Hoekstra, A. Babul, D. D. Balam, and P. L. Capak, *Astrophys.J.* **668**, 806 (2007), 0706.3048.
80. H. D. Falcke, M. P. van Haarlem, A. G. de Bruyn, R. Braun, H. J. A. Röttgering, B. Stappers, W. H. W. M. Boland, H. R. Butcher, E. J. de Geus, L. V. Koopmans, *et al.*, *Highlights of Astronomy* **14**, 386 (2007), arXiv:astro-ph/0610652.
81. K. Barbary, K. S. Dawson, K. Tokita, G. Aldering, R. Amanullah, N. V. Connolly, M. Doi, L. Faccioli, V. Fadeyev, A. S. Fruchter, *et al.*, *Astrophys.J.* **690**, 1358 (2009), 0809.1648.
82. N. Soker, A. Frankowski, and A. Kashi, *ArXiv e-prints* (2008), 0812.1402.
83. A. Kusenko, *Nuclear Physics B Proceedings Supplements* **173**, 24 (2007), arXiv:astro-ph/0609375.
84. D. Gorbunov, A. Khmelnsky, and V. Rubakov, *Journal of High Energy Physics* **12**, 55 (2008),

0805.2836.

85. D. Gorbunov, A. Khmelnitsky, and V. Rubakov, *Journal of Cosmology and Astro-Particle Physics* **10**, 41 (2008), 0808.3910.
86. D. Boyanovsky, *Phys. Rev. D* **78**, 103505 (2008), 0807.0646.
87. A. Badertscher, P. Crivelli, W. Fetscher, U. Gendotti, S. N. Gninenko, V. Postoev, A. Rubbia, V. Samoylenko, and D. Sillou, *Phys. Rev. D* **75**, 032004 (2007), arXiv:hep-ex/0609059.
88. P. Ciarcelluti, *International Journal of Modern Physics D* **14**, 187 (2005), arXiv:astro-ph/0409630.
89. P. Ciarcelluti, *International Journal of Modern Physics D* **14**, 223 (2005), arXiv:astro-ph/0409633.
90. S. I. Blinnikov, in *Radio Universe at Ultimate Angular Resolution* (2008), URL <http://radioastron.ru/index2.php?engdep=12&meet=3>.